

## Review

# Homocysteine, folic acid and vitamin B<sub>12</sub> in relation to pre- and postnatal health aspects

Rima Obeid and Wolfgang Herrmann\*

Department of Clinical Chemistry, Saarland Medical School, Homburg, Germany

### Abstract

Studies linking hyperhomocysteinemia (HHCY) and B-vitamin deficiency to some health aspects in children have been accumulating. Low B-vitamin status in early life, even as early as the time of conception, may endanger the potential for new life and may negatively influence the health of the offspring. Early abortion, pregnancy complications and poor pregnancy outcomes have been linked to elevated concentrations of total plasma homocysteine (tHcy) and low folate or vitamin B<sub>12</sub>. Maternal vitamin status predicts that of fetuses and neonates. Lactating women are likely to experience low micronutrient status, which might affect breast-milk composition and hence the nutritional status of their breast-fed infants. Elevated concentrations of methylmalonic acid (MMA) is common in infants (age <6 months), which may indicate a transient inadequate vitamin B<sub>12</sub> status. Deficiency of B-vitamins might confer deleterious effects on the physical and mental health of the child, such as impaired growth, gross motor function, poor school performance and other adaptive skills. The importance of maintaining adequate B-vitamin status during periods of progressive growth and development should be emphasized because symptoms related to folate and vitamin B<sub>12</sub> deficiency are difficult to detect. Serum levels of tHcy and MMA should be estimated in several target groups of children, pregnant and lactating women and those planning for pregnancy. Concentrations of tHcy and MMA are useful indicators of B-vitamin status in the pediatric laboratory. Using these functional markers may facilitate detecting sub-optimal B-vitamin status in children.

**Keywords:** folate; homocysteine; pediatrics; vitamin B<sub>12</sub>.

### Homocysteine: a new marker in pediatric settings

Numerous studies have investigated concentrations of plasma total homocysteine (tHcy) in children dur-

ing the last few years. Concentrations of tHcy and methylmalonic acid (MMA) are sensitive and specific metabolic markers for B-vitamin status (1, 2). The classic hematological manifestations associated with folate or vitamin B<sub>12</sub> deficiency are not uniquely expressed (3). Moreover, serum concentrations of these vitamins have poor specificity and sensitivity (2). Therefore, the assessment of tHcy and MMA offers a useful diagnostic tool in the pediatric laboratory.

Concentrations of tHcy in infants (range 4–9 µmol/L) are relatively higher than in older children (1). Vitamin B<sub>12</sub> is the major modulator of tHcy in infants younger than 1 year and folate becomes a main determinant of tHcy at older age (1). In children over 1 year, as well as in adults, age becomes a significant factor that modulates tHcy levels (1). In general, children (>1 year) have lower tHcy concentrations than adolescents (4). Boys have slightly higher tHcy than girls and this sexual dimorphism of tHcy concentrations occurs as early as 10 years of age (5). Elevated tHcy concentration in children may be related to inherited or acquired factors (6) (Table 1).

### Homocysteine, a risk factor for pregnancy complication and outcome

The importance of B-vitamins in human reproduction has been repeatedly addressed. Low maternal status of B-vitamins or hyperhomocysteinemia (HHCY) has been linked to pregnancy complications and poor birth outcomes (7–9).

Concentrations of tHcy decrease during pregnancy and are usually lower in pregnant women supplemented with folic acid compared to unsupplemented women (10). An elevated concentration of tHcy during pregnancy was independently associated with the risk of recurrent pregnancy loss (7), preeclampsia, prematurity and low birth weight (8–10). The association between maternal HHcy and intrauterine growth retardation has not been confirmed by all investigators (11). Different findings from studies in Europe and Canada could be related to different study designs or to improved folate status in Canada since the fortification of food with folic acid in 1998. In line with this suggestion, Canadian women homozygous for the mutation in the methylenetetrahydrofolate reductase gene (*MTHFR* C677T) who were not taking multivitamins during pregnancy were at increased risk of having a child with intrauterine growth restriction (12). The importance of vitamin B<sub>6</sub> during pregnancy has also been outlined in several studies (13).

\*Corresponding author: Prof. Dr. Wolfgang Herrmann, Department of Clinical Chemistry and Laboratory Medicine, School of Medicine, Saarland University, Gebäude 57 Annexbau, 66421 Homburg, Germany  
Phone: +49-6841-1630700, Fax: +49-6841-1630703,  
E-mail: kcwher@uniklinik-saarland.de

**Table 1** Common causes of hyperhomocysteinemia in newborns and children.

Inherited
• Hetero- or homozygosity for mutations in genes encoding enzymes (e.g., CBS or MS)
• <i>MTHFR</i> mutation
• Cbl E/G and F defects
• Inherited transcobalamin deficiency
Acquired
Primary
• Folate, vitamin B <sub>12</sub> or B <sub>6</sub> deficiency
• Vegetarian, vegan or macrobiotic diet
• Poverty or malnutrition
• Newborns of vitamin-deficient mothers
Secondary
• HIV-infected children
• Malabsorption, bacterial overgrowth
• Medications (anti-epileptics, theophylline)
• Renal dysfunction

CBS, cystathione- $\beta$ -synthase; MS, methionine synthase; Cbl, cobalamin.

Poor maternal vitamin B<sub>6</sub> status during pregnancy has been related to low birth weight of the offspring (13).

Taken together, it is recommended that women of reproductive age should increase their dietary intake of folate, vitamin B<sub>12</sub> and B<sub>6</sub>. Deficiency of one of these vitamins should be suspected in women with a current or a previous unfavorable event during pregnancy.

### Homocysteine in diseased children

Homocysteinuria, the disease firstly linked to disturbed tHcy metabolism, is a rare genetic defect that causes a marked reduction in the activity of cystathionine- $\beta$ -synthetase and therefore severe HHcy (tHcy > 100  $\mu$ mol/L). HHcy is a risk factor for ocular, cerebral, skeletal and clotting pathologies in homocysteinuric children (14). Large-scale screening programs of newborns are not currently recommended (15). However, the importance of selective screening for HHcy in pediatric patients has recently been highlighted (16). In contrast to severe HHcy, modest elevation of tHcy (10–30  $\mu$ mol/L) is frequently encountered. Nevertheless, the implications of HHcy in children have not been fully explored.

Children treated with certain medications, such as anti-epileptic drugs, had elevated concentrations of tHcy (17). Mild to moderate HHcy is a risk factor for occlusive vascular disease, cerebrovascular lesions and thrombosis in children (17–19). HHcy is also very common in pediatric renal patients, but can be lowered by therapeutic doses of folic acid (20). Evidence available from vitamin intervention studies in adult renal patients is promising and suggests that similar studies should be implemented in children as well.

Higher concentrations of tHcy and lower concentrations of folate and vitamin B<sub>12</sub> were found in HIV-infected children (21). Malabsorption, low dietary intake and increased demands may contribute to micronutrient deficiency in this case. Neurological and neurodevelopmental disorders are common in such children (21). B-vitamins are required for the pro-

duction of S-adenosylmethionine, the methyl donor for numerous biochemical reactions, including the formation of neurotransmitters. Therefore, B-vitamin deficiency may cause deprivation of the methyl groups in the central nervous system.

A notable exception of elevated tHcy concentrations in the context of childhood disease is the case of Down syndrome (trisomy 21). The gene encoding cystathionine- $\beta$ -synthetase enzyme located on chromosome 21 is overexpressed in children with Down syndrome. Accordingly, concentrations of tHcy, methionine and S-adenosylmethionine were lower in children with Down syndrome compared to normal siblings (22).

Vitamin treatment lowered tHcy concentrations and improved some measures of endothelial function in pediatric patients (23). Prospective studies may clarify whether tHcy-lowering therapy may improve the progression and outcome of tHcy-related morbidities in children.

### Periods of high demands for B-vitamins

Folate and vitamin B<sub>12</sub> play a fundamental role in the biosynthesis of purine and pyrimidine and in maintaining DNA methylation. A high rate of DNA synthesis and gene expression characterizes all rapidly dividing cells. Therefore, increased demands for B-vitamins in pregnant women and during early life are expected.

Body stores of folates are limited and depletion occurs rapidly. A marginal, usually asymptomatic, low vitamin status during the periconception period may traverse into a deficiency state during pregnancy or lactation. Prolonged lactation may deplete serum folate and cause tHcy elevation in mothers with a low daily intake ( $\leq 380$   $\mu$ g) (24). Nevertheless, prolonged lactation did not cause folate depletion in mothers taking 1 mg of folic acid daily (24). Other target groups for which adequate intake and status of the B-vitamins should be stressed are pre-school and school-age children (25). Long-term deficiency may have deleterious neurological and developmental consequences (discussed below) (25, 26).

### Predictors of tHcy and B-vitamin status in early life

Maternal concentrations of vitamin B<sub>12</sub> and folate predict those in the neonate at birth (27, 28). The placenta sequesters vitamin B<sub>12</sub>, B<sub>6</sub> and folate. Moreover, the concentrations of these vitamins are two- to six-fold higher in cord blood than in maternal serum (27, 28). This may support the high metabolic rate in the developing fetus.

Fetal and maternal tHcy concentrations are strongly correlated (27–29). Concentrations of tHcy at the time of labor were lower in mothers taking folic acid compared to unsupplemented mothers (10). Low concentration of vitamin B<sub>12</sub> (or high MMA) in mothers was associated with lower B<sub>12</sub> and higher MMA in their newborns (30). Moreover, babies who had the highest

MMA (and lowest B<sub>12</sub>) values at birth also had the highest MMA values (and lowest B<sub>12</sub>) at 6 weeks (30).

Concentrations of tHcy in neonates are influenced by nutritional factors (folate, B<sub>12</sub> and B<sub>6</sub>) (31–34). Vitamin B<sub>12</sub> intake may be limited during the first year of life because of a low vitamin content in breast milk and an immature intrinsic factor system. The estimated stores of vitamin concentrations B<sub>12</sub> in the neonatal liver are expected to cover the requirements for this vitamin during the first year of life. Vitamin stores in newborns of depleted mothers may be readily depleted, especially in breast-fed infants. Previous studies demonstrated that breast-fed infants of vegetarian mothers excrete more MMA in their urine than infants of non-vegetarian women (32, 34). Moreover, lower concentrations of tHcy and higher concentrations of B<sub>12</sub> were found in formula-fed infants compared to breast-fed infants (35). Several studies reported a positive correlation between maternal serum and milk content of the vitamin (32, 33). Moreover, the occurrence and the reversible nature of methylmalonic aciduria in breast-fed infants from deficient mothers confirm that maternal B<sub>12</sub> status determines that of the newborns (32, 34, 35). Vitamin B<sub>12</sub> status cannot be easily restored in newborns of deficient mothers, especially if vitamin intake continues to be in the lower range (36, 37). As a consequence, newborns of deficient mothers may develop a real deficiency syndrome later in life (36–38).

### Occurrence of a sub-normal B-vitamin status in early life

A high incidence of B-vitamin deficiency has been documented worldwide. This occurs frequently in the context of poverty or prolonged inadequate intake. Vitamin B<sub>12</sub> deficiency is highly prevalent in vegetarians, macrobiotic communities and in some developing countries (33, 36).

A transient inadequate status of vitamin B<sub>12</sub> may be common at age <6 months (1). Concentrations of MMA and tHcy were higher in infants (<6 months) than that in older children (1). Measurement of MMA in conjunction with tHcy improved the specificity of these tests (1). The usefulness of serum levels of holotranscobalamin, the vitamin B<sub>12</sub> portion bound to transcobalamin, in estimating vitamin B<sub>12</sub> status in infants should be determined.

Vitamin B<sub>12</sub> deficiency was very prevalent in Guatemalan lactating mothers and their infants at 3 months postpartum (33). B<sub>12</sub> content in breast milk was low in 31% of these women and correlated to urinary MMA in their infants (33). An intake below the recommended daily amount (RDA) for B<sub>12</sub> (0.3 µg/day) was recorded in 16% of the infants (33). In a study of lactating women in Mexico, dietary intake of vitamin B<sub>12</sub> was only 50% of that recommended and 62% of the mothers had a low content of B<sub>12</sub> in their breast milk (39). Low vitamin B<sub>12</sub> status was also common in Guatemalan and Kenyan schoolchildren (40, 41). Furthermore, folate deficiency was very common (43%) and folate intake was also below recom-

mended during lactation (500 µg/day). In addition, folate content in human milk was not adequate to support the requirements of many infants (39, 42).

A high incidence of B<sub>12</sub> deficiency was found in infants consuming a macrobiotic diet (36). Ingesting a strict vegetarian diet in early life led to a low vitamin B<sub>12</sub> status in adolescence, despite increasing B<sub>12</sub> intake since the age of 6 years (43).

### Homocysteine and B-vitamins in relation to birth defects

Several congenital anomalies may be prevented by folic acid supplementation before and during early pregnancy (44). The exact mechanism by which folic acid exerts its protective effect is unclear. However, the role of the vitamin in one-carbon metabolism, nucleic acid synthesis and the integrity of DNA are plausible explanations.

Neural tube defects (NTDs) are the most common congenital malformations that have been linked to maternal folate deficiency and HHCY (44–46). In addition, concentrations of tHcy were higher in mothers of children with NTDs compared to mothers of normal children (45). Concentrations of folate were lower in children with spina bifida compared to normal children (46). Low blood folate was associated with an odds ratio of 2.6 for having NTD and of 3.1 for being a mother of a child with an NTD (46). Recent strategies for the primary prevention of NTDs include preconceptional folic acid supplementation (47). In 1998, the US Food and Drug Administration mandated that all grain products be fortified with folic acid (140 µg per 100 g of grain). A similar policy has also been put in place in Canada, Chile and Hungary. This national health policy is proposed to provide a daily intake of at least 400 µg/day folic acid in women of reproductive age. Folate status and concentrations of tHcy have been improved in women of childbearing age after such fortification (48). This metabolic improvement has also been associated with a marked reduction in birth defects (between 15% and 30%) (49).

Down syndrome or trisomy 21 is a common genetic disorder that is due to disturbed segregation of chromosome 21 during meiosis. DNA hypomethylation, due to methyl-group deficiency, influences chromosome segregation. Folate and vitamin B<sub>12</sub> are required for controlled DNA methylation and genomic stability. The effect of abnormal folate metabolism on the maternal risk of having a child with Down syndrome has been examined (50). A significant increase in tHcy levels was observed in mothers of children with Down syndrome and an approximately three-fold higher risk was found in mothers homozygous for the *MTHFR* C677T mutation compared to mothers without the T substitution (50).

Genes encoding enzymes that participate in the distribution of folate forms or in folate metabolism and transport are important for embryonic development. Several mutations in genes involved in folate and methionine metabolism have been linked to the risk

**Table 2** Target groups who should increase their intake of vitamin B<sub>12</sub> and folic acid.

- Pregnant and lactating women
- Mothers with previous pregnancy loss, complication or poor pregnancy outcome
- Women planning for pregnancy
- Infants of strict vegetarian mothers
- Infants with unexplained neurological syndromes
- Infants with developmental delay
- Infants with siblings affected by homocysteinuria, birth defects or developmental delay
- Breast-fed infants of vitamin-deficient mothers
- Pediatric patients (renal patients, diabetes, epilepsy)

of some congenital anomalies (50–54). A specific pattern of gene-nutrient interaction has been described. Therefore, improvement of the maternal nutritional status may modify the associated risk of certain genetic variants (46, 51). Subjects homozygous for the *MTHFR* C677T mutation have higher tHcy levels when folate and/or B<sub>12</sub> supply is limited. A positive association has been reported between the *MTHFR* TT genotype and NTDs (55). The risk for NTDs increased in *MTHFR* carriers who had low folate levels (46). Other mutations in genes encoding homocysteine remethylation enzymes may also increase the risk of NTDs (54).

### Role of vitamin B<sub>12</sub> status in postnatal development

Deficiency of one or more of the B-vitamins may lead to irreversible neurological disorders. Folate or vitamin B<sub>12</sub> deficiencies occur mostly in infants of deficient mothers (33, 34, 36, 38, 56, 57). Symptoms of vitamin B<sub>12</sub> deficiency in neonates become evident at 4–9 months of age (38). A serious neurological syndrome and developmental disorders have been described in breast-fed infants of strict vegetarian mothers (38, 58). A few studies relating B<sub>12</sub> status to cognitive development in children are available. A significant association was found between biochemical markers of B<sub>12</sub> status and performance on tests measuring cognitive function in children (59, 60). School achievement was negatively related to vitamin B<sub>12</sub> status in Guatemalan children (25, 60). Moreover, lower scores in some measures of cognitive performance were found in adolescents who previously consumed a macrobiotic diet compared to those who consumed an omnivorous diet from birth onwards (37). Vitamin B<sub>12</sub> deficiency in infants was associated with marked developmental regression, poor brain growth or poor intellectual outcome (25, 60). Other signs of developmental delay include impaired communicative reactions, and fine and gross motor functions.

In conclusion, deficiency of the B-vitamins in infancy leads to long-term metabolic and neurological abnormalities. Table 2 summarizes target groups for which the intake of folate and vitamin B<sub>12</sub> should be increased. Early recognition and prevention of these deficiencies should be emphasized. Age-specific ref-

erence ranges for blood concentrations of folate, B<sub>12</sub> and their surrogate markers should be established, because adults reference values cannot be extrapolated to infants and children.

### References

1. Bjorke Monsen AL, Refsum H, Markestad T, Ueland PM. Cobalamin status and its biochemical markers methylmalonic acid and homocysteine in different age groups from 4 days to 19 years. *Clin Chem* 2003;49:2067–75.
2. Savage DG, Lindenbaum J, Stabler SP, Allen RH. Sensitivity of serum methylmalonic acid and total homocysteine determinations for diagnosing cobalamin and folate deficiencies. *Am J Med* 1994;96:239–46.
3. Obeid R, Geisel J, Schorr H, Hubner U, Herrmann W. The impact of vegetarianism on some haematological parameters. *Eur J Haematol* 2002;69:275–9.
4. Delvin EE, Rozen R, Merouani A, Genest J Jr, Lambert M. Influence of methylenetetrahydrofolate reductase genotype, age, vitamin B-12, and folate status on plasma homocysteine in children. *Am J Clin Nutr* 2000;72:1469–73.
5. Must A, Jacques PF, Rogers G, Rosenberg IH, Selhub J. Serum total homocysteine concentrations in children and adolescents: results from the third National Health and Nutrition Examination Survey (NHANES III). *J Nutr* 2003;133:2643–9.
6. Rosenblatt DS, Whitehead VM. Cobalamin and folate deficiency: acquired and hereditary disorders in children. *Semin Hematol* 1999;36:19–34.
7. Nelen WL. Hyperhomocysteinaemia and human reproduction. *Clin Chem Lab Med* 2001;39:758–63.
8. Ronnenberg AG, Goldman MB, Chen D, Aitken IW, Willett WC, Selhub J, et al. Preconception homocysteine and B vitamin status and birth outcomes in Chinese women. *Am J Clin Nutr* 2002;76:1385–91.
9. Vollset SE, Refsum H, Irgens LM, Emblem BM, Tverdal A, Gjessing HK, et al. Plasma total homocysteine, pregnancy complications, and adverse pregnancy outcomes: the Hordaland Homocysteine Study. *Am J Clin Nutr* 2000;71:962–8.
10. Murphy MM, Scott JM, Arijia V, Molloy AM, Fernandez-Ballart JD. Fetal homocysteine and birth weight are affected by maternal homocysteine before conception and throughout pregnancy. *Clin Chem* 2004;50:1406–12.
11. Infante-Rivard C, Rivard GE, Gauthier R, Theoret Y. Unexpected relationship between plasma homocysteine and intrauterine growth restriction. *Clin Chem* 2003;49:1476–82.
12. Infante-Rivard C, Rivard GE, Yotov WV, Genin E, Guiguet M, Weinberg C, et al. Absence of association of thrombophilia polymorphisms with intrauterine growth restriction. *N Engl J Med* 2002;347:19–25.
13. Reinken L, Dapunt O. Vitamin B6 nutrition during pregnancy. *Int J Vitam Nutr Res* 1978;48:341–7.
14. McCully KS. Vascular pathology of homocysteinemia: implications for the pathogenesis of arteriosclerosis. *Am J Pathol* 1969;56:111–28.
15. Refsum H, Grindflek AW, Ueland PM, Fredriksen A, Meyer K, Ulvik A, et al. Screening for serum total homocysteine in newborn children. *Clin Chem* 2004;50:1769–84.
16. Vilaseca MA, Moyano D, Artuch R, Ferrer I, Pineda M, Cardo E, et al. Selective screening for hyperhomocysteinemia in pediatric patients. *Clin Chem* 1998;44:662–4.
17. Cardo E, Vilaseca MA, Campistol J, Artuch R, Colome C, Pineda M. Evaluation of hyperhomocysteinemia in children with stroke. *Eur J Paediatr Neurol* 1999;3:113–7.



18. Hogeveen M, Blom HJ, Van Amerongen M, Boogmans B, Van Beynum IM, Van De Bor M. Hyperhomocysteinemia as risk factor for ischemic and hemorrhagic stroke in newborn infants. *J Pediatr* 2002;141:429–31.
19. Kosch A, Koch HG, Heinecke A, Kurnik K, Heller C, Nowak-Gottl U, et al. Increased fasting total homocysteine plasma levels as a risk factor for thromboembolism in children. *Thromb Haemost* 2004;91:308–14.
20. Kang HG, Lee BS, Hahn H, Lee JH, Ha IS, Cheong HI, et al. Reduction of plasma homocysteine by folic acid in children with chronic renal failure. *Pediatr Nephrol* 2002;17:511–4.
21. Boudes P, Zittoun J, Sobel A. Folate, vitamin B<sub>12</sub>, and HIV infection. *Lancet* 1990;335:1401–2.
22. Pogribna M, Melnyk S, Pogribny I, Chango A, Yi P, James SJ. Homocysteine metabolism in children with Down syndrome: in vitro modulation. *Am J Hum Genet* 2001;69:88–95.
23. Pena AS, Wiltshire E, Gent R, Hirte C, Couper J. Folic acid improves endothelial function in children and adolescents with type 1 diabetes. *J Pediatr* 2004;144:500–4.
24. Mackey AD, Picciano MF. Maternal folate status during extended lactation and the effect of supplemental folic acid. *Am J Clin Nutr* 1999;69:285–92.
25. Penland JG, Allen LH, Boy E, DeBaessa Y, Rogers LM. Adaptive functioning, behavior problems and school performance of Guatemalan school children with deficient, marginal and normal plasma B-12 [abstract]. *FASEB J* 2000;14:A561.
26. Lindenbaum J, Healton EB, Savage DG, Brust JC, Garrett TJ, Podell ER, et al. Neuropsychiatric disorders caused by cobalamin deficiency in the absence of anemia or macrocytosis. *N Engl J Med* 1988;318:1720–8.
27. Obeid R, Munz W, Jäger M, Schmidt W, Herrmann W. Biochemical indexes of the B-vitamins in cord serum are predicted by maternal B-vitamin status. *Am J Clin Nutr* 2005;82:133–9.
28. Guerra-Shinohara EM, Morita OE, Peres S, Pagliusi RA, Sampaio Neto LF, D'Almeida V, et al. Low ratio of S-adenosylmethionine to S-adenosylhomocysteine is associated with vitamin deficiency in Brazilian pregnant women and newborns. *Am J Clin Nutr* 2004;80:1312–21.
29. Malinow MR, Rajkovic A, Duell PB, Hess DL, Upson BM. The relationship between maternal and neonatal umbilical cord plasma homocyst(e)ine suggests a potential role for maternal homocyst(e)ine in fetal metabolism. *Am J Obstet Gynecol* 1998;178:228–33.
30. Bjorke Monsen AL, Ueland PM, Vollset SE, Guttormsen AB, Markestad T, Solheim E, et al. Determinants of cobalamin status in newborns. *Pediatrics* 2001;108:624–30.
31. Minet JC, Bisse E, Aebischer CP, Beil A, Wieland H, Lutschg J. Assessment of vitamin B-12, folate, and vitamin B-6 status and relation to sulfur amino acid metabolism in neonates. *Am J Clin Nutr* 2000;72:751–7.
32. Specker BL, Black A, Allen L, Morrow F. Vitamin B-12: low milk concentrations are related to low serum concentrations in vegetarian women and to methylmalonic aciduria in their infants. *Am J Clin Nutr* 1990;52:1073–6.
33. Casterline JE, Allen LH, Ruel MT. Vitamin B-12 deficiency is very prevalent in lactating Guatemalan women and their infants at three months postpartum. *J Nutr* 1997;127:1966–72.
34. Specker BL, Miller D, Norman EJ, Greene H, Hayes KC. Increased urinary methylmalonic acid excretion in breast-fed infants of vegetarian mothers and identification of an acceptable dietary source of vitamin B-12. *Am J Clin Nutr* 1988;47:89–92.
35. Specker BL, Brazier W, Ho ML, Norman EJ. Urinary methylmalonic acid excretion in infants fed formula or human milk. *Am J Clin Nutr* 1990;51:209–11.
36. Schneede J, Dagnelie PC, van Staveren WA, Vollset SE, Refsum H, Ueland PM. Methylmalonic acid and homocysteine in plasma as indicators of functional cobalamin deficiency in infants on macrobiotic diets. *Pediatr Res* 1994;36:194–201.
37. Louwman MW, van Dusseldorp M, van de Vijver FJ, Thomas CM, Schneede J, Ueland PM, et al. Signs of impaired cognitive function in adolescents with marginal cobalamin status. *Am J Clin Nutr* 2000;72:762–9.
38. Higginbottom MC, Sweetman L, Nyhan WL. A syndrome of methylmalonic aciduria, homocystinuria, megaloblastic anemia and neurologic abnormalities in a vitamin B12-deficient breast-fed infant of a strict vegetarian. *N Engl J Med* 1978;299:317–23.
39. Black AK, Allen LH, Pelto GH, de Mata MP, Chavez A. Iron, vitamin B-12 and folate status in Mexico: associated factors in men and women and during pregnancy and lactation. *J Nutr* 1994;124:1179–88.
40. Rogers LM, Boy E, Miller JW, Green R, Sabel JC, Allen LH. High prevalence of cobalamin deficiency in Guatemalan schoolchildren: associations with low plasma holotranscobalamin II and elevated serum methylmalonic acid and plasma homocysteine concentrations. *Am J Clin Nutr* 2003;77:433–40.
41. Siekmann JH, Allen LH, Bwibo NO, Demment MW, Murphy SP, Neumann CG. Kenyan school children have multiple micronutrient deficiencies, but increased plasma vitamin B-12 is the only detectable micronutrient response to meat or milk supplementation. *J Nutr* 2003;133:3972S–80S.
42. Villalpando S, Latulippe ME, Rosas G, Irurita MJ, Picciano MF, O'Connor DL. Milk folate but not milk iron concentrations may be inadequate for some infants in a rural farming community in San Mateo, Capulhuac, Mexico. *Am J Clin Nutr* 2003;78:782–9.
43. van Dusseldorp M, Schneede J, Refsum H, Ueland PM, Thomas CM, de Boer E, et al. Risk of persistent cobalamin deficiency in adolescents fed a macrobiotic diet in early life. *Am J Clin Nutr* 1999;69:664–71.
44. Czeizel AE, Dudas I. Prevention of the first occurrence of neural-tube defects by periconceptional vitamin supplementation. *N Engl J Med* 1992;327:1832–5.
45. Steegers-Theunissen RP, Boers GH, Trijbels FJ, Finkelstein JD, Blom HJ, Thomas CM, et al. Maternal hyperhomocysteinemia: a risk factor for neural-tube defects? *Metabolism* 1994;43:1475–80.
46. Christensen B, Arbour L, Tran P, Leclerc D, Sabbaghian N, Platt R, et al. Genetic polymorphisms in methylenetetrahydrofolate reductase and methionine synthase, folate levels in red blood cells, and risk of neural tube defects. *Am J Med Genet* 1999;84:151–7.
47. Czeizel AE. Primary prevention of neural-tube defects and some other major congenital abnormalities: recommendations for the appropriate use of folic acid during pregnancy. *Paediatr Drugs* 2000;2:437–49.
48. Ray JG, Vermeulen MJ, Boss SC, Cole DE. Increased red cell folate concentrations in women of reproductive age after Canadian folic acid food fortification. *Epidemiology* 2002;13:238–40.
49. Williams LJ, Mai CT, Edmonds LD, Shaw GM, Kirby RS, Hobbs CA, et al. Prevalence of spina bifida and anencephaly during the transition to mandatory folic acid fortification in the United States. *Teratology* 2002;66:33–9.
50. James SJ, Pogribna M, Pogribny IP, Melnyk S, Hine RJ, Gibson JB, et al. Abnormal folate metabolism and mutation in the methylenetetrahydrofolate reductase gene may be maternal risk factors for Down syndrome. *Am J Clin Nutr* 1999;70:495–501.
51. Molloy AM, Mills JL, Kirke PN, Ramsbottom D, McPartlin JM, Burke H, et al. Low blood folates in NTD pregnancies are only partly explained by thermolabile 5,10-methyle-

- netetrahydrofolate reductase: low folate status alone may be the critical factor. *Am J Med Genet* 1998;78: 155–9.
52. Wilson A, Platt R, Wu Q, Leclerc D, Christensen B, Yang H, et al. A common variant in methionine synthase reductase combined with low cobalamin (vitamin B12) increases risk for spina bifida. *Mol Genet Metab* 1999;67: 317–23.
53. Chango A, Emery-Fillon N, de Courcy GP, Lambert D, Pfister M, Rosenblatt DS, et al. A polymorphism (80G → A) in the reduced folate carrier gene and its associations with folate status and homocysteinemia. *Mol Genet Metab* 2000;70:310–5.
54. Zhu H, Wicker NJ, Shaw GM, Lammer EJ, Hendricks K, Suarez L, et al. Homocysteine remethylation enzyme polymorphisms and increased risks for neural tube defects. *Mol Genet Metab* 2003;78:216–21.
55. van der Put NM, Eskes TK, Blom HJ. Is the common 677C → T mutation in the methylenetetrahydrofolate reductase gene a risk factor for neural tube defects? A meta-analysis. *QJM* 1997;90:111–5.
56. Sklar R. Nutritional vitamin B12 deficiency in a breast-fed infant of a vegan-diet mother. *Clin Pediatr (Phila)* 1986;25:219–21.
57. VanderJagt DJ, Spelman K, Ambe J, Datta P, Blackwell W, Crossey M, et al. Folate and vitamin B12 status of adolescent girls in northern Nigeria. *J Natl Med Assoc* 2000;92:334–40.
58. Kuhne T, Bubl R, Baumgartner R. Maternal vegan diet causing a serious infantile neurological disorder due to vitamin B12 deficiency. *Eur J Pediatr* 1991;150:205–8.
59. Graham SM, Arvela OM, Wise GA. Long-term neurologic consequences of nutritional vitamin B12 deficiency in infants. *J Pediatr* 1992;121:710–4.
60. Allen LH, Penland JG, Boy E, DeBaessa Y, Rogers LM. Cognitive and neuromotor performance of Guatemalan schoolers with deficient, marginal, and normal plasma vitamin B-12 [abstract]. *FASEB J* 1999;13:A544.